

Charge and spin correlations of a Peierls insulator after a quench

Martin Hohenadler

Institut für Theoretische Physik und Astrophysik, Universität Würzburg, 97074 Würzburg, Germany

(Dated: March 25, 2013)

Electron-phonon coupling plays a central role for time-dependent phenomena in condensed matter, for example in photo-excitation experiments. We use the continuous-time quantum Monte Carlo method to study the real-time evolution of charge and spin correlation functions of a Peierls insulator after a quench to a noninteracting Hamiltonian. This approach gives exact results, and fully takes into account quantum phonon effects without relying on a Hilbert space truncation. It is also free from a dynamical sign problem. Since thermalization is absent for a quench to an integrable model, we observe dephasing over a time scale set by the electronic hopping integral. Our exact results provide a benchmark for more realistic calculations, and may be directly applicable to experiments with cold atoms or trapped ions.

PACS numbers: 71.38.-k, 71.45.Lr, 63.20.kd

I. INTRODUCTION

The enormous current interest in the time evolution or nonequilibrium behavior of quantum systems is driven by significant developments in the field of experimental physics. Of particular importance are time-resolved studies of cold atoms in optical lattices following the seminal work by Greiner *et al.*,¹ and the progress with femtosecond spectroscopy,^{2–5} including time-resolved photoemission spectroscopy.^{6,7} These methods enable researchers to carry out pump-probe experiments and to study photo-excitation and photo-induced phase transitions.^{4,8–14}

Theoretical studies are particularly difficult for quantum systems with strong correlations, as relevant for cold atoms and several classes of materials. Whereas much progress has been made in developing methods suitable to study correlated systems in equilibrium, the description of time-dependent or nonequilibrium problems is extremely challenging. As a result, it is currently not possible to reliably describe experiments. Promising directions of research include time-dependent density matrix renormalization group methods,¹⁵ extensions of quantum cluster methods such as dynamical mean-field theory^{16,17} and the variational cluster approach,¹⁸ as well as extensions of Luttinger liquid theory.¹⁹ A review of theoretical methods for correlated systems has been given by Eckstein *et al.*²⁰

Whereas these ideas have successfully been applied to models of interacting spins, fermions or bosons, much less is known about phenomena related to electron-phonon interaction. The latter plays a crucial role since phonons are often the fastest channel available for thermalization via scattering.²¹ However, the infinite bosonic Hilbert space as well as the resulting retarded electron-electron interaction pose a particular challenge to theoretical methods. Whereas exact results have been obtained for a single polaron,^{22–24} a single bipolaron,²⁵ and a single-molecule device,²⁶ lattice models with finite band fillings are much more challenging. Recent work along these lines makes use of the density matrix renormalization group on small systems,²⁷ the so-called double-phonon cloud

method,²⁸ restricted phonon Hilbert spaces,²⁹ as well as classical phonons.³⁰

A particularly interesting class of experiments involves photo-induced phase transition across a metal-insulator Peierls transition.^{4,8–14,31} A laser pulse excites the system out of the insulating, charge-ordered state, as visible for example from the time dependence of the gap in the band structure (probed by time and angle-resolved photoemission spectroscopy)¹³ or from the reflectivity.¹¹ After the photo-induced melting of charge order, the system relaxes back towards its original state. Such experiments also reveal the typically very different timescales for electron and ion dynamics. So far, a photo-induced phase transition of a Peierls system has been considered theoretically in Ref. 29 using a classical phonon approximation. Since nonlocal correlations are a defining feature of charge ordered states, local approximations such as dynamical-mean field theory are of limited use.

Motivated by the ongoing effort to improve the available theoretical tools for the study of electron-phonon systems, we consider here the quantum quench from a Peierls insulator to noninteracting fermions by switching off the electron-phonon interaction. The model we use is the one-dimensional Holstein model. The time evolution of charge and spin correlation functions after the quench is studied with an extension of the continuous-time quantum Monte Carlo method.³² Although such a quench does not reflect the complex experimental situations encountered in pump-probe experiments, it provides us with a rare opportunity to obtain exact results for the time evolution of an electron-phonon system. Our findings can serve as a nontrivial benchmark for future numerical calculations (for example with methods that rely on a truncation of the phonon Hilbert space), and may even be relevant for experimental realizations of electron-phonon models in optical lattices^{33–36} or ion traps.^{37,38}

The paper is organized as follows. In Sec. II we discuss the model, in Sec. III we provide details about the method used, in Sec. IV we present our results, and Sec. V contains our conclusions.

II. MODEL

The Peierls metal-insulator transition in one dimension can be investigated in the framework of Holstein's molecular crystal model,³⁹ defined by the Hamiltonian

$$\hat{H} = \sum_{k\sigma} \epsilon(k) c_{k\sigma}^\dagger c_{k\sigma} + \sum_i \left(\frac{\hat{P}_i^2}{2M} + \frac{K\hat{Q}_i^2}{2} \right) - g(t) \sum_i \hat{Q}_i (\hat{n}_i - 1). \quad (1)$$

The first term, referred to as $\hat{H}_0$ in the following, describes nearest-neighbor hopping of electrons with the usual tight-binding band structure $\epsilon(k) = -2J \cos(k)$; we have set the lattice constant equal to one, and take the hopping integral as the energy unit. The second and third terms correspond to the lattice degrees of freedom and the electron-phonon coupling. The lattice is described by means of harmonic oscillators with frequency $\omega_0 = \sqrt{K/M}$, displacement $\hat{Q}_i$ and momentum $\hat{P}_i$, and the coupling is of the density-displacement type proposed by Holstein;³⁹ $g(t)$ is the coupling constant. We use the usual definitions $\hat{n}_{i\sigma} = c_{i\sigma}^\dagger c_{i\sigma}$, $\hat{n}_i = \sum_\sigma \hat{n}_{i\sigma}$, and $n = \langle \hat{n}_i \rangle$. A useful dimensionless coupling parameter is given by the ratio $\lambda = 2\varepsilon_p/W$, where $\varepsilon_p = g^2/2K$ is the polaron binding energy, and $W = 4J$ is the bare bandwidth. We consider a half filled band, *i.e.*, $\langle \hat{n}_i \rangle = 1$.

As discussed in Sec. I, we study a quantum quench to a noninteracting Hamiltonian. This is achieved by a time-dependent coupling constant $g(t)$. Explicitly, we take

$$g(t) = \begin{cases} g, & t = 0, \\ 0, & t > 0. \end{cases} \quad (2)$$

Because there is no coupling between electrons and the lattice after such a quench, the time evolution will be determined by the hopping term $\hat{H}_0$ only.

III. METHOD

The continuous-time quantum Monte Carlo (CTQMC) method is based on a weak-coupling Dyson expansion of the path integral for the partition function.³² For finite systems, all possible vertex contributions can be summed stochastically, thereby giving exact results with only statistical errors. The CTQMC method has been used quite extensively to solve impurity problems in the context of dynamical mean-field theory, see Ref. 40 for a review, as well as for lattice simulations.

Because the method is formulated in terms of an action rather than a Hamiltonian, it permits us to simulate models with retarded interactions. Such interactions naturally arise as a result of electron-phonon interaction, and models such as Eq. (1) can be simulated by integrating out the lattice degrees of freedom analytically.⁴¹ This

approach has been applied to study spinless⁴² and spinful Holstein^{43,44} and Holstein-Hubbard models,⁴⁵ as well as models with a nonlocal electron-phonon interaction.⁴⁶ Numerical details can be found in Ref. 45. Because the method relies on an analytical integration over phonon degrees of freedom, lattice properties such as local displacements cannot be directly measured.

Quantum Monte Carlo approaches to time-dependent problems are inherently limited by the dynamical sign problem related to the time evolution operator $\exp(i\hat{H}t)$. Nevertheless, progress has been made in applying such methods to simplified situations where the sign problem can be avoided or limited, including quantum quenches to noninteracting systems,⁴⁷ and studies of (effective) impurity problems. For the CTQMC method used here, the first application to a nonequilibrium problem involving phonons was achieved in Ref. 48, and a review of previous work is found in Ref. 40. Recently, the weak-coupling CTQMC method was formulated on the Keldysh contour.⁴⁷

In principle, the method of Ref. 47 can be extended to include a retarded electron-electron interaction on the Keldysh contour. The resulting algorithm would permit us to study rather general problems, but with the usual limitation to short times originating from the dynamical sign problem.⁴⁷ Instead, we follow a simpler route sufficient to study quenches to noninteracting systems, based on the idea used by Goth and Assaad in the context of the auxiliary-field quantum Monte Carlo method.⁴⁷ If we are only interested in the time evolution *after* the electron-phonon coupling g has been switched off [see Eq. (2)], interaction vertices are restricted to imaginary times on the Keldysh contour. The time dependence of observables is then fully determined by the noninteracting Hamiltonian $\hat{H}_0$, the first term in Eq. (1).

The numerical method proceeds as follows. The interacting Holstein model (1) is simulated with the CTQMC method using standard Monte Carlo updates (addition and removal of vertices, flipping of Ising spins⁴¹) to obtain configurations reflecting the initial, correlated state at time $t = 0$. The interacting single-particle Green's function matrix at equal imaginary times,

$$[\mathbf{G}_\sigma(t=0)]_{ij} = \langle c_{i\sigma} c_{j\sigma}^\dagger \rangle, \quad (3)$$

is calculated for a given vertex configuration from the Dyson equation.^{32,41} From $\mathbf{G}_\sigma(t=0)$, all relevant correlation functions and observables at $t = 0$ can be obtained using Wick's theorem.^{32,41} For quenches of the form (2), expectation values at times $t > 0$ follow from the time-propagated Green function

$$\mathbf{G}_\sigma(t) = e^{-i\hat{H}_0 t} \mathbf{G}_\sigma(t=0) e^{i\hat{H}_0 t}, \quad (4)$$

in combination with Wick's theorem. The additional numerical effort compared to a time-independent simulation amounts to the evaluation of Eq. (4) and of observables of interest for each value of t . Results for each t are averaged over different Monte Carlo configurations. Whereas

equal-time observables can easily be calculated in this way, time-displaced correlation functions require a formulation on the Keldysh contour.

IV. RESULTS

For a half-filled band, the Holstein model (1) describes a transition from a metal to a Peierls insulator driven by the electron-phonon interaction. At $T = 0$, the Peierls state has long-range charge order with wavevector $q = 2k_F$, and a gap in the single-particle spectrum.⁴⁹ In addition, a spin gap presumably exists for any nonzero electron-phonon interaction, as a result of electrons pairing into singlets.⁴⁵ Whereas the model is dimerized and insulating for any $\lambda > 0$ in the mean-field limit $\omega_0 = 0$,⁴⁹ numerical results^{50–53} indicate a nonzero critical value for $\omega_0 > 0$ as a result of quantum lattice fluctuations. For an overview of previous work, see Ref. 45.

Here, we consider parameters for which Hamiltonian (1) is in the insulating Peierls phase. Explicitly, we take $\lambda = 0.5$, and different values of the phonon frequency, $\omega_0 = 0.5J$, and $5J$. According to quantum Monte Carlo results,⁵³ $\lambda = 0.5$ is well inside the ordered phase for $\omega_0 = 0.5J$ (the critical coupling is about 0.25), and just above the critical coupling for $\omega_0 = 5J$. Since our simulations are carried out at a low but finite temperature (we consider a fixed system size $L = 22$ and $\beta J = 2L = 44$), the system will be in a thermal initial state before the quench. However, the results shown below are practically identical for $\beta J = L$ and $\beta J = 2L$.

Among the equal-time observables accessible with the present method, the real-space charge and spin correlation functions

$$\begin{aligned} S_\rho(r) &= \langle \hat{n}_r \hat{n}_0 \rangle, \\ S_\sigma(r) &= \langle \hat{S}_r^z \hat{S}_0^z \rangle, \end{aligned} \quad (5)$$

are of particular interest. Since we consider a quench to the noninteracting Hamiltonian, we compare the correlation functions to those of free fermions. For the latter, Luttinger liquid theory gives⁵⁴

$$S_\rho(r) = S_\sigma(r) = -\frac{1}{(\pi r)^2} + \frac{C}{r^2} \cos(2k_F r). \quad (6)$$

Here, we have set $K_\rho = K_\sigma = 1$, as appropriate for free fermions. In contrast, the Peierls state is formally described by $K_\rho = 0$ (implying long-range $2k_F$ charge order) and $K_\sigma = 0$ (reflecting the spin gap which leads to exponentially suppressed spin correlations).⁴⁵

Figure 1 shows results for $S_\rho(r, t)$ for $\omega_0 = 0.5J$ and $\lambda = 0.5$, as calculated from the time-dependent Green's function (4). Panels (a)–(d) correspond to different times t in the interval $[0, 0.7J]$, whereas panel (e) shows a closeup. Full lines show the charge correlation function at time t after the quench, whereas dotted lines are results for free fermions.




FIG. 1. (Color online) Charge correlations $S_\rho(r, t)$ (full lines) at different times t after the quench. Here, $\lambda = 0.5$, $\omega_0 = 0.5J$, $L = 22$ and $\beta J = 44$. Dotted lines correspond to $S_\rho(r)$ for $\lambda = 0$. Panel (e) shows a closeup of panel (d), and additional results for $t = 0.63J$.

At $t = 0$, Fig. 1(a), we observe the clear $2k_F$ charge correlations expected for the Peierls state. At $t > 0$, after the quench to $\lambda = 0$, these oscillations decay over a time scale of about $J/\sqrt{2}$, after which $2k_F$ oscillations are practically absent at large distances. Shortly before that time, namely for $t \approx 0.63J$, small-amplitude $2k_F$ oscillations comparable to the free-fermion results are visible, see Fig. 1(e). For $t > 0.7J$, a revival towards a charge-ordered state begins. Whereas the charge correlations for $t = 0.7J$ in Fig. 1(d) agree well with those for the noninteracting system on the scale of the figure, the closeup in Fig. 1(e) reveals that noticeable differences remain at small distances even for the ‘optimal time’ of $t = 0.63J$.

Figure 2 shows the corresponding results for the spin correlation function $S_\sigma(r, t)$. The times shown are the same as in Fig. 1. In contrast to $S_\rho(r, t)$, Fig. 1(a), there are no discernible $2k_F$ correlations at $t = 0$, in accordance with the nonzero spin gap of the Peierls state. After the quench, the differences from the noninteracting results visible at short distances diminish as a function of time. For $t = 0.7J$, similar to the charge correlations, $S_\sigma(r, t)$ looks very similar to the noninteracting result on the scale of Fig. 2(d), but the closeup in Fig. 2(e) again reveals clear deviations at small distances. Interestingly,




FIG. 2. (Color online) Spin correlations $S_\sigma(r, t)$ (full lines) at different times t after the quench. Here, $\lambda = 0.5$, $\omega_0 = 0.5J$, $L = 22$ and $\beta J = 44$. Dotted lines correspond to $S_\sigma(r)$ for $\lambda = 0$. Panel (e) shows a closeup of panel (d), and additional results for $t = 0.63J$.

the results for $S_\sigma(r, t = 0.7J)$ in Fig. 2(e) agree within the symbol size used with those for $S_\rho(r, t = 0.7J)$, despite substantial differences at $t = 0$.

Because the system is quenched to a noninteracting and hence integrable Hamiltonian ($\hat{H}_0$), real thermalization is not possible.⁵⁵ Instead, the time evolution seen in Figs. 1 and 2 is related to dephasing. Indeed, we observe the expected periodic but incomplete revivals of the system at longer times, see Fig. 3 and discussion below. Conserved quantities such as the total energy and the momentum distribution function are independent of t . Since the evolution is determined by $\hat{H}_0$ [cf. Eq. (4)], there is only a single energy scale, namely the hopping integral J . However, we can modify the initial state by varying the parameters λ and ω_0 . As emphasized before, our method fully accounts for quantum phonon fluctuations in the initial state. An increase of the phonon frequency brings the initial state much closer to the Peierls phase boundary.⁵³ To illustrate the dependence of charge order and its time evolution on the phonon frequency, we show in Fig. 3 the double occupancy, $\langle n_{i\uparrow}n_{i\downarrow} \rangle$, which reflects the pairing of electrons at the same site in the Peierls state, and its decay after the quench. Comparing the cases of $\omega_0 = 0.5J$ and $\omega_0 = 5J$, Fig. 3 reveals a signif-




FIG. 3. (Color online) Double occupation as a function of time for $\lambda = 0.5$ and two values of the phonon frequency. Here, $L = 22$ and $\beta J = 44$. The vertical line illustrates the noninteracting value 0.25.

icantly smaller double occupation at $t = 0$ for $\omega_0 = 5J$ than for $\omega_0 = 0.5J$. At $t > 0$, we observe a fast decay of the double occupation, with periodic oscillations around the noninteracting value of 0.25. The time scale associated with this decay agrees with the decay of $2k_F$ charge correlations in Fig. 1. A partial revival to about half the value at $t = 0$ is visible at time $t \approx 5J$, and reappears with this period also at later times (data not shown).

V. CONCLUSIONS

Motivated by an increasing number of experiments on ultrafast dynamics after photo-induced phase transitions, and by the enormous challenge of numerically describing time-dependent phenomena in systems with electron-phonon interaction, we considered the quench of a Peierls insulator to a noninteracting Hamiltonian. Using an extension of the continuous-time quantum Monte Carlo method, we presented exact results for equal-time charge and spin correlation functions. The time evolution of the latter is related to dephasing, and is determined by the only remaining energy scale after the quench, namely the hopping J . We find that the dominant $2k_F$ charge correlations of the initial Peierls state decay over a time scale of about $J/\sqrt{2}$. After that time, we find that spin and charge correlations are virtually identical.

In the present case, the electron-phonon coupling and the quantum nature of phonons only play a role for the initial state. Increasing (decreasing) the phonon frequency (the coupling) moves the system closer to the metallic state, thereby suppressing charge order. Importantly, the results shown cannot be obtained by considering the adiabatic limit $\omega_0 \rightarrow \infty$, where the Holstein model maps onto an attractive Hubbard model. The reason is that the attractive Hubbard model at half filling always remains metallic, with the Luttinger liquid pa-

parameter K_p fixed to one by symmetry.⁵⁶

The exact results presented here can serve as benchmarks for more realistic calculations using, for example, the density-matrix renormalization group. In the latter, the phonon Hilbert space has to be truncated for the initial state as well as for the time evolution. Finally, our findings may even become directly relevant for experiments with cold atoms or trapped ions; several different proposals exist how to realize electron-phonon Hamiltonians in such systems.^{33–35,37,38}

ACKNOWLEDGMENTS

Computer time at the Jülich Supercomputing Centre, support from the DFG Grant No. Ho 4489/2-1 (Forschergruppe FOR 1807), and discussions with F. F. Assaad, F. Goth and M. Sentef are gratefully acknowledged.

-
- ¹ M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, *Nature* (London) **415**, 39 (2002)
 - ² A. Pashkin, M. Porer, M. Beyer, K. W. Kim, A. Dubroka, C. Bernhard, X. Yao, Y. Dagan, R. Hackl, A. Erb, J. Demsar, R. Huber, and A. Leitenstorfer, *Phys. Rev. Lett.* **105**, 067001 (2010)
 - ³ C. Gadermaier, A. S. Alexandrov, V. V. Kabanov, P. Kusar, T. Mertelj, X. Yao, C. Manzoni, D. Brida, G. Cerullo, and D. Mihailovic, *Phys. Rev. Lett.* **105**, 257001 (2010)
 - ⁴ Y. Kawakami, T. Fukatsu, Y. Sakurai, H. Unno, H. Itoh, S. Iwai, T. Sasaki, K. Yamamoto, K. Yakushi, and K. Yonemitsu, *Phys. Rev. Lett.* **105**, 246402 (2010)
 - ⁵ S. Wall, D. Brida, S. R. Clark, H. P. Ehrke, D. Jaksch, A. Ardavan, S. Bonora, H. Uemura, Y. Takahashi, T. Hasegawa, H. Okamoto, G. Cerullo, and A. Cavalleri, *Nature Phys.* **7**, 114 (2011)
 - ⁶ L. Perfetti, P. A. Loukakos, M. Lisowski, U. Bovensiepen, H. Eisaki, and M. Wolf, *Phys. Rev. Lett.* **99**, 197001 (2007)
 - ⁷ R. Cortés, L. Rettig, Y. Yoshida, H. Eisaki, M. Wolf, and U. Bovensiepen, *Phys. Rev. Lett.* **107**, 097002 (2011)
 - ⁸ M. Chollet, L. Guerin, N. Uchida, S. Fukaya, H. Shimoda, T. Ishikawa, K. Matsuda, T. Hasegawa, A. Ota, H. Yamochi, G. Saito, R. Tazaki, S.-i. Adachi, and S.-y. Koshihara, *Science* **307**, 86 (2005)
 - ⁹ S. Iwai, K. Yamamoto, A. Kashiwazaki, F. Hiramatsu, H. Nakaya, Y. Kawakami, K. Yakushi, H. Okamoto, H. Mori, and Y. Nishio, *Phys. Rev. Lett.* **98**, 097402 (2007)
 - ¹⁰ C. Kübler, H. Ehrke, R. Huber, R. Lopez, A. Halabica, R. F. Haglund, and A. Leitenstorfer, *Phys. Rev. Lett.* **99**, 116401 (2007)
 - ¹¹ A. Tomeljak, H. Schäfer, D. Städter, M. Beyer, K. Biljakovic, and J. Demsar, *Phys. Rev. Lett.* **102**, 066404 (2009)
 - ¹² S. Hellmann, M. Beye, C. Sohrt, T. Rohwer, F. Sorgenfrei, H. Redlin, M. Kalläne, M. Marczyński-Bühlöw, F. Hennies, M. Bauer, A. Föhlisch, L. Kipp, W. Wurth, and K. Rossnagel, *arXiv:1004.4790*(2010)
 - ¹³ R. G. Moore, V. Brouet, R. He, D. H. Lu, N. Ru, J.-H. Chu, I. R. Fisher, and Z.-X. Shen, *Phys. Rev. B* **81**, 073102 (2010)
 - ¹⁴ S. Miyashita, Y. Tanaka, S. Iwai, and K. Yonemitsu, *J. Phys. Soc. Jpn.* **79**, 034708 (2010)
 - ¹⁵ U. Schollwöck, *Annals of Physics* **326**, 96 (2011)
 - ¹⁶ P. Schmidt and H. Monien, *cond-mat/0202046*(2002)
 - ¹⁷ J. K. Freericks, V. M. Turkowski, and V. Zlatić, *Phys. Rev. Lett.* **97**, 266408 (2006)
 - ¹⁸ M. Knap, W. von der Linden, and E. Arrigoni, *Phys. Rev. B* **84**, 115145 (2011)
 - ¹⁹ C. Karrasch, J. Rentrop, D. Schuricht, and V. Meden, *Phys. Rev. Lett.* **109**, 126406 (2012)
 - ²⁰ M. Eckstein, A. Hackl, S. Kehrein, M. Kollar, M. Moeckel, P. Werner, and F. Wolf, *The European Physical Journal Special Topics* **180**, 217 (2009)
 - ²¹ M. Sentef, A. F. Kemper, B. Moritz, J. K. Freericks, Z.-X. Shen, and T. P. Devereaux, *arXiv:1212.4841*(2012)
 - ²² D. Golež, J. Bonča, L. Vidmar, and S. A. Trugman, *Phys. Rev. Lett.* **109**, 236402 (2012)
 - ²³ L.-C. Ku and S. A. Trugman, *Phys. Rev. B* **75**, 014307 (2007)
 - ²⁴ H. Fehske, G. Wellein, and A. R. Bishop, *Phys. Rev. B* **83**, 075104 (2011)
 - ²⁵ D. Golež, J. Bonča, and L. Vidmar, *Phys. Rev. B* **85**, 144304 (2012)
 - ²⁶ Y. Vinkler, A. Schiller, and N. Andrei, *arXiv:1109.5590*(2011)
 - ²⁷ H. Matsueda, S. Sota, T. Tohyama, and S. Maekawa, *J. Phys. Soc. Jpn.* **81**, 013701 (2012)
 - ²⁸ G. De Filippis, V. Cataudella, E. A. Nowadnick, T. P. Devereaux, A. S. Mishchenko, and N. Nagaosa, *Phys. Rev. Lett.* **109**, 176402 (2012)
 - ²⁹ J. D. Lee, P. Moon, and M. Hase, *Phys. Rev. B* **84**, 195109 (2011)
 - ³⁰ K. Yonemitsu and N. Maeshima, *Phys. Rev. B* **79**, 125118 (2009)
 - ³¹ F. Schmitt, P. S. Kirchmann, U. Bovensiepen, R. G. Moore, L. Rettig, M. Krenz, J.-H. Chu, N. Ru, L. Perfetti, D. H. Lu, M. Wolf, I. R. Fisher, and Z.-X. Shen, *Science* **321**, 1649 (2008)
 - ³² A. N. Rubtsov, V. V. Savkin, and A. I. Lichtenstein, *Phys. Rev. B* **72**, 035122 (2005)
 - ³³ F. Herrera and R. V. Krems, *Phys. Rev. A* **84**, 051401 (2011)
 - ³⁴ J. P. Hague and C. MacCormick, *New Journal of Physics* **14**, 033019 (2012)
 - ³⁵ E. Pazy and A. Vardi, *Phys. Rev. A* **72**, 033609 (2005)
 - ³⁶ M. Atala, M. Aidelsburger, J. T. Barreiro, D. Abanin, T. Kitagawa, E. Demler, and I. Bloch, *arXiv:1212.0572*(2012)
 - ³⁷ A. Mezzacapo, J. Casanova, L. Lamata, and E. Solano, *Phys. Rev. Lett.* **109**, 200501 (2012)
 - ³⁸ V. M. Stojanović, T. Shi, C. Bruder, and J. I. Cirac, *Phys. Rev. Lett.* **109**, 250501 (2012)
 - ³⁹ T. Holstein, *Ann. Phys. (N.Y.)* **8**, 325; **8**, 343 (1959)
 - ⁴⁰ E. Gull, A. J. Millis, A. I. Lichtenstein, A. N. Rubtsov, M. Troyer, and P. Werner, *Rev. Mod. Phys.* **83**, 349 (2011)
 - ⁴¹ F. F. Assaad and T. C. Lang, *Phys. Rev. B* **76**, 035116 (2007)

- ⁴² M. Hohenadler, H. Fehske, and F. F. Assaad, Phys. Rev. B **83**, 115105 (2011)
- ⁴³ F. F. Assaad, Phys. Rev. B **78**, 155124 (2008)
- ⁴⁴ M. Hohenadler and F. F. Assaad, J. Phys.: Condens. Matter **25**, 014005 (2013)
- ⁴⁵ M. Hohenadler and F. F. Assaad, Phys. Rev. B **87**, 075149 (2013)
- ⁴⁶ M. Hohenadler, F. F. Assaad, and H. Fehske, Phys. Rev. Lett. **109**, 116407 (2012)
- ⁴⁷ F. Goth and F. F. Assaad, Phys. Rev. B **85**, 085129 (2012)
- ⁴⁸ L. Mühlbacher and E. Rabani, Phys. Rev. Lett. **100**, 176403 (2008)
- ⁴⁹ J. E. Hirsch and E. Fradkin, Phys. Rev. B **27**, 4302 (1983)
- ⁵⁰ E. Jeckelmann, C. Zhang, and S. R. White, Phys. Rev. B **60**, 7950 (1999)
- ⁵¹ H. Fehske, G. Hager, and E. Jeckelmann, Europhys. Lett. **84**, 57001 (2008)
- ⁵² R. T. Clay and R. P. Hardikar, Phys. Rev. Lett. **95**, 096401 (2005)
- ⁵³ R. P. Hardikar and R. T. Clay, Phys. Rev. B **75**, 245103 (2007)
- ⁵⁴ H. J. Schulz, Phys. Rev. Lett. **64**, 2831 (1990)
- ⁵⁵ M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, Phys. Rev. Lett. **98**, 050405 (2007)
- ⁵⁶ T. Giamarchi, *Quantum physics in one dimension* (Clarendon Press, Oxford, 2004)